


Anaphylaxis Emergency Plan: _____ (name)

This person has a potentially life-threatening allergy (anaphylaxis) to:



PHOTO

(Check the appropriate boxes.)

Food(s): _____

Insect stings
 Other: _____

Epinephrine Auto-Injector: Expiry Date: _____ / _____

Dosage:

EpiPen® Jr. 0.15 mg EpiPen® 0.30 mg

Location of Auto-Injector(s): _____

Previous anaphylactic reaction: Person is at greater risk.

Asthmatic: Person is at greater risk. If person is having a reaction and has difficulty breathing, give epinephrine auto-injector before asthma medication.

A person having an anaphylactic reaction might have ANY of these signs and symptoms:

- **Skin system:** hives, swelling (face, lips, tongue), itching, warmth, redness
- **Respiratory system (breathing):** coughing, wheezing, shortness of breath, chest pain or tightness, throat tightness, hoarse voice, nasal congestion or hay fever-like symptoms (runny, itchy nose and watery eyes, sneezing), trouble swallowing
- **Gastrointestinal system (stomach):** nausea, pain or cramps, vomiting, diarrhea
- **Cardiovascular system (heart):** paler than normal skin colour/blue colour, weak pulse, passing out, dizziness or lightheadedness, shock
- **Other:** anxiety, sense of doom (the feeling that something bad is about to happen), headache, uterine cramps, metallic taste

Early recognition of symptoms and immediate treatment could save a person's life.

Act quickly. The first signs of a reaction can be mild, but symptoms can get worse very quickly.

1. **Give epinephrine auto-injector** (e.g. EpiPen®) **at the first sign of a known or suspected anaphylactic reaction.** (See attached instruction sheet.)
2. **Call 9-1-1** or local emergency medical services. Tell them someone is having a life-threatening allergic reaction.
3. **Give a second dose of epinephrine as early as 5 minutes after the first dose if there is no improvement in symptoms.**
4. **Go to the nearest hospital immediately (ideally by ambulance),** even if symptoms are mild or have stopped. The reaction could worsen or come back, even after proper treatment. Stay in the hospital for an appropriate period of observation as decided by the emergency department physician (generally about 4-6 hours).
5. **Call emergency contact person (e.g. parent, guardian).**

Emergency Contact Information

Name	Relationship	Home Phone	Work Phone	Cell Phone

The undersigned patient, parent, or guardian authorizes any adult to administer epinephrine to the above-named person in the event of an anaphylactic reaction, as described above. This protocol has been recommended by the patient's physician.

Patient/Parent/Guardian Signature

Date

Physician Signature On file

Date



Blue to the sky. Orange to the thigh.

How to use EpiPen® and EpiPen® Jr (epinephrine) Auto-injectors.

Remove the EpiPen® Auto-Injector from the carrier tube and follow these 2 simple steps:



- Hold firmly with orange tip pointing downward.
- Remove blue safety cap by pulling straight up.
Do not bend or twist.



- Swing and push orange tip firmly into mid-outer thigh until you hear a 'click'.
- Hold on thigh for several seconds.



Built-in needle protection

- After injection, the orange cover automatically extends to ensure the needle is never exposed.



After using EpiPen®, you must seek immediate medical attention or go to the emergency room. For the next 48 hours, you must stay close to a healthcare facility or be able to call 911.

For more information visit the consumer site EpiPen.ca.

EpiPen® and EpiPen® Jr (epinephrine) Auto-Injectors are indicated for the emergency treatment of anaphylactic reactions in patients who are determined to be at increased risk for anaphylaxis, including individuals with a history of anaphylactic reactions. Selection of the appropriate dosage strength is determined according to patient body weight.

EpiPen® and EpiPen® Jr Auto-Injectors are designed as emergency supportive therapy only. They are not a replacement for subsequent medical or hospital care. After administration, patients should seek medical attention immediately or go to the emergency room. For the next 48 hours, patients must stay within close proximity to a healthcare facility or where they can call 911. To ensure this product is right for you, always read and follow the label. Please consult the Consumer Information leaflet in your product package for complete dosage and administration instructions.



© 2015 Pfizer Canada Inc., Kirkland, Quebec H9J 2M5 • Toll free: 1-877-EPIPEN1 (1-877-374-7361)
EpiPen®, EpiPen® Jr are registered trademarks of Mylan, Inc. licensed exclusively to its wholly-owned affiliate,
Mylan Specialty, L.P.; sub-licensee, Pfizer Canada Inc., Kirkland, Quebec H9J 2M5
D00005059



Trusted for over 25 years.